

Exact Simulation for a Class of Tempered Stable and Related Distributions

ANGELOS DASSIOS and YAN QU, London School of Economics
HONGBIAO ZHAO, Shanghai University of Finance and Economics

In this article, we develop a new scheme of exact simulation for a class of tempered stable (TS) and other related distributions with similar Laplace transforms. We discover some interesting integral representations for the underlying density functions that imply a unique simulation framework based on a backward recursive procedure. Therefore, the foundation of this simulation design is very different from existing schemes in the literature. It works pretty efficiently for some subclasses of TS distributions, where even the conventional acceptance-rejection mechanism can be avoided. It can also generate some other distributions beyond the TS family. For applications, this scheme could be easily adopted to generate a variety of TS-constructed random variables and TS-driven stochastic processes for modelling observational series in practice. Numerical experiments and tests are performed to demonstrate the accuracy and effectiveness of our scheme.

CCS Concepts: • **Mathematics of computing** → **Random number generation**;

Additional Key Words and Phrases: Monte carlo simulation, exact simulation, backward recursive scheme, stable distribution, tempered stable distribution, exponentially tilted stable distribution, Lévy process, Lévy subordinator, leptokurtosis

ACM Reference format:

Angelos Dassios, Yan Qu, and Hongbiao Zhao. 2018. Exact Simulation for a Class of Tempered Stable and Related Distributions. *ACM Trans. Model. Comput. Simul.* 28, 3, Article 20 (July 2018), 21 pages.
<https://doi.org/10.1145/3184453>

1 INTRODUCTION

From the empirical analysis, it is well known that asset returns in financial markets are not normally distributed, and the observations often present skewness and heavy tails, especially during crises. Stable distributions, first introduced by Lévy (1925), is a very important class that often provides an alternative model for price changes observed in financial markets. More precisely, the distributions form a four-parameter family of *infinitely divisible distributions* (Sato 1999, p. 31), which can be defined by the logarithm of the characteristic function,

The corresponding author Hongbiao Zhao would like to acknowledge the financial support from the *National Natural Science Foundation of China* (No. 71401147) and the research fund provided by Shanghai University of Finance and Economics.

Authors' addresses: A. Dassios and Y. Qu, Department of Statistics, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom; emails: {a.dassios, y.qu3}@lse.ac.uk; H. Zhao (corresponding author), School of Statistics and Management, Shanghai University of Finance and Economics, and Shanghai Institute of International Finance and Economics, No. 777 Guoding Road, Shanghai 200433, China; email: h.zhao1@lse.ac.uk.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2018 ACM 1049-3301/2018/07-ART20 \$15.00

<https://doi.org/10.1145/3184453>

$$\ln \phi(v) = \begin{cases} i\ell v - \gamma^\alpha |v|^\alpha \left(1 - i\zeta \tan\left(\frac{\alpha}{2}\pi\right) \text{sign}(v)\right), & \alpha \neq 1, \\ i\ell v - \gamma |v| \left(1 + i\zeta \frac{2}{\pi} \ln |v|\right), & \alpha = 1, \end{cases} \quad v \in \mathbb{R},$$

where $\phi(v)$ is the characteristic function of stable distributions, i is the imaginary unit, i.e., $i^2 = -1$, $\alpha \in (0, 2]$ is the *stability index*, $\zeta \in [-1, 1]$ is the *skewness parameter*, $\gamma \in \mathbb{R}^+$ is the *scale parameter*, and $\ell \in \mathbb{R}$ is the *location parameter*. Their great flexibility attracted many scholars in economics and finance at least since the 1960s, and the study was initialised by the seminal work of Mandelbrot (1960, 1961, 1963a, 1963b) and followed by Fama (1965) and Fama and Roll (1968, 1971). Tremendous applications of stable distributions and their various extensions can be found in the literature (e.g., option pricing and portfolio management); see a detailed survey by McCulloch (1996).

However, stable distributions could involve infinite moments, and this property limits its applications. For instance, this model may lead to an infinite price in the option pricing; see Carr and Wu (2003). Therefore, the distribution, in particular its tail, needs to be adjusted. *Tempered stable* (TS), or *exponentially tilted stable* (ETS) distribution¹ provides a very important class that attempts to overcome this limitation of the original stable distribution. It was initially proposed by Tweedie (1984); see also Hougaard (1986) and Jorgensen (1987). An additional parameter of *tilt* is included so that their moments exist. The TS law is a natural generalisation, as it inherits desirable features from the stable distribution, such as infinite divisibility, skewness, and leptokurtosis. The resulting tails are lighter than those of the stable but heavier than those of the Gaussian, so it is more suitable for modelling financial returns. Moreover, for those distributions that are obtained by tilting *one-sided stable distributions*² with $\zeta = 1$, namely the *positive TS distributions*, they can be further adopted as the building blocks to construct various stochastic processes (such as Lévy TS subordinators and processes) that facilitate numerous applications in finance and many other fields, e.g., for modelling the dynamics of stock prices, stochastic volatilities, or interest rates just to name a few; see Barndorff-Nielsen and Shephard (2001a, 2002), Schoutens (2003), Cont and Tankov (2004), Kyprianou (2006), and, more recently, Barndorff-Nielsen et al. (2012).

Still, the challenging issue is that, with only certain exceptions, density functions of stable and TS distributions have no explicit expressions (Zolotarev 1986, p. 2). Hence, simulation-based methods, in particular exact simulation schemes without bias, become extremely important in practice, especially for the purposes of statistical inference, numerical validation, risk analysis, and pricing exotic financial derivatives.

Let us first have a brief review of some important simulation schemes for stable, TS, and their related distributions in the literature. For a stable distribution in general, the standard algorithm for exact simulation is the classical method of Chambers et al. (1976) (CMS),³ which is the extension of the work by Kanter (1975). This state-of-the-art algorithm is remarkably elegant, accurate, and fast, since the procedure is *direct*, which means that even the acceptance-rejection (A/R) procedure is not required. However, for the TS distribution, it is usually hard to find such a *direct* scheme for exact simulation. The very popular algorithm is the classical *simple stable rejection* (SSR) scheme (with the algorithm provided in Appendix A). It is straightforward to be implemented, as it is developed by simply combining the classical CMS method with the A/R scheme; see Brix (1999) and Baeumer and Meerschaert (2010) for more details. More recently, Devroye (2009) developed

¹Here, we adopt the naming for the TS distribution similarly as Küchler and Tappe (2013) and Kim and Kim (2016). In fact, the class of TS distributions has been recently enlarged by Rosiński (2007) and Grabchak et al. (2012) in a variety of specifications, so we consider them as *generalised TS distributions*.

²The stable distributions are one-sided when $\alpha \in (0, 1)$ and $\zeta = \pm 1$.

³The detailed proof is given by Weron (1996).

an alternative algorithm of *double rejection (DR)* method such that the complexity is uniformly bounded. Hofert (2011a, 2011b) suggested a *fast rejection* algorithm to enhance the original SSR scheme. All the three algorithms are *exact*⁴ and based on the A/R scheme, which can be applied to the class of positive TS distributions with stability index $\alpha \in (0, 1)$ ⁵.

For the existing algorithms to simulate a TS random variable (RV), each of them has its own advantages but also have some limitations for some choices of parameters. It is therefore very useful to develop some alternative simulation schemes to sample the TS family. In this article, we discover a multiple integral representation for the density functions of TS distributions with stability index in a general *dyadic form* $\frac{q}{2^n} \in (0, 1)$ for $q, n \in \mathbb{N}^+$, the integral representation allows us to develop an alternative simulation framework for the TS distributions with stability index $\frac{q}{2^n} \in (0, 1)$. In particular, when q is a binary number, an extremely efficient *direct* scheme can be established, which means that the algorithm only involves the explicit inverse transformation without the A/R mechanism. On one hand, this alternative algorithm could be very helpful for numerical validating for TS-based stochastic models, because a variety of choices of q, n combined with the different values of parameters β and θ would provide enough flexibility for the purpose of numerical testing, and at the same time, the corresponding computational cost of implementing TS-based models could be substantially reduced. On the other hand, since the foundation of our simulation design is indeed very different from existing schemes in the literature, this representation therefore allows us to design the simulation algorithms for a wider range of distributions with Lévy measures beyond the general TS family.

The article is organised as follows: In Section 2, we first present the fundamental results on the multiple integral representation for a special recursive type of Laplace transform functions and provide an simulation framework for some distributions beyond the TS $\frac{q}{2^n}$ -family. In Section 3, we derive a new simulation scheme, namely the *backward recursive (BR) scheme*, for exact simulation based on the multiple integral representation for the density of $\frac{q}{2^n}$ -TS distributions. In addition, we outline the improved algorithms tailored for two important subclasses, one is a *direct* scheme for the $\frac{1}{2^n}$ -family, and the other is an enhanced scheme for the $\frac{3}{2^n}$ -family. In Section 4, numerical experiments for our algorithms as well as the associated comparisons with other schemes have been carried out and reported. Section 5 makes a conclusion of this article, and proposes some issues for possible further extensions and future research.

2 FUNDAMENTAL RESULTS

Before investigating the TS distributions, let us first introduce a general integral representation for a special type of Laplace transform functions based on the explicit formula for the inverse Gaussian (IG) density. This special type of Laplace transform has an interesting recursive structure, which characterises a variety of different distributions even beyond the TS family. The integral representation leads to an recursive simulation scheme if the initial distribution is given. The theoretical foundation in general is illustrated in Theorem 2.1 below.

THEOREM 2.1. *If a series of non-negative functions $\hat{f}_1(v), \dots, \hat{f}_n(v)$ for any $n \in \mathbb{N}^+$ possess a recursive structure of*

$$\hat{f}_k(v) = \begin{cases} \hat{f}_1(v), & k = 1, \\ a_k \sqrt{b_k + c_k v + \hat{f}_{k-1}(v)} - a_k \sqrt{b_k}, & k = 2, \dots, n, \end{cases} \quad (1)$$

⁴On the other hand, some approximation-based algorithms have also been proposed in the literature, see Bondesson (1982), Rosiński (2001), and Ridout (2009), however, they obviously involve numerical errors.

⁵For the class of $\alpha \in (1, 2)$, a non-exact simulation algorithm based on the A/R scheme with Gaussian approximation is developed in Kawai and Masuda (2011).

where $a_k > 0$ and $b_k, c_k \geq 0$, then, $e^{-\hat{f}_n(v)}$ can be represented by

$$e^{-\hat{f}_n(v)} = \begin{cases} \int_0^\infty e^{-vs_1} f_1(s_1) ds_1, & n = 1, \\ \int_0^\infty \cdots \int_0^\infty e^{-s_1 v - \sum_{k=2}^n c_k s_k v} f_1(s_1 | s_2) \frac{a_2 s_3 e^{-\frac{(\sqrt{2b_2 s_2 - \frac{a_2 s_3}{\sqrt{2}}})^2}{2s_2}}}{2\sqrt{\pi s_2^3}} \times \cdots \times \frac{a_n e^{-\frac{(\sqrt{2b_n s_n - \frac{a_n}{\sqrt{2}}})^2}{2s_n}}}{2\sqrt{\pi s_n^3}} ds_n \cdots ds_1, & n > 1, \end{cases} \quad (2)$$

where $f_1(s_1) = \mathcal{L}_{s_1}^{-1}\{e^{-\hat{f}_1(v)}\}$ and $f_1(s_1 | s_2) = \mathcal{L}_{s_1}^{-1}\{e^{-s_2 \hat{f}_1(v)}\}$.

PROOF. We use $IG(\frac{a}{b}, a^2)$ to denote an IG distribution⁶ with mean $\frac{a}{b}$ and the shape parameter a^2 where $a > 0, b \geq 0$. It is well known that its Laplace transform is given by

$$\mathbb{E} \left[e^{-vIG(\frac{a}{b}, a^2)} \right] = e^{-a(\sqrt{b^2 + 2v} - b)}.$$

Since Equation (2) holds for $n = 2$. The proof for the general case can be conducted by the mathematical induction. Assuming Equation (2) also holds for an arbitrary integer $n = j > 2$, given $s_{j+1} \in (0, \infty)$, we have

$$\begin{aligned} e^{-\hat{f}_j(v)s_{j+1}} &= \exp \left(-a_j s_{j+1} \left[\sqrt{b_j + c_j v + \hat{f}_{j-1}(v)} - \sqrt{b_j} \right] \right) \\ &= \int_0^\infty \cdots \int_0^\infty e^{-\left(s_1 + \sum_{k=2}^j c_k s_k\right)v} f_1(s_1 | s_2) \frac{a_2 s_3 e^{-\frac{(\sqrt{2b_2 s_2 - \frac{a_2 s_3}{\sqrt{2}}})^2}{2s_2}}}{2\sqrt{\pi s_2^3}} \times \cdots \times \frac{a_j s_{j+1} e^{-\frac{(\sqrt{2b_j s_j - \frac{a_j s_{j+1}}{\sqrt{2}}})^2}{2s_j}}}{2\sqrt{\pi s_j^3}} ds_j \cdots ds_1. \end{aligned}$$

Then, for $n = j + 1$, we have

$$\begin{aligned} e^{-\hat{f}_{j+1}(v)} &= \int_0^\infty e^{-c_{j+1} s_{j+1} v} e^{-\hat{f}_j(v)s_{j+1}} \frac{a_{j+1}}{2\sqrt{\pi s_{j+1}^3}} e^{-\frac{(\sqrt{2b_{j+1} s_{j+1} - \frac{a_{j+1}}{\sqrt{2}}})^2}{2s_{j+1}}} ds_{j+1} \\ &= \int_0^\infty \cdots \int_0^\infty e^{-\left(s_1 + \sum_{k=2}^{j+1} c_k s_k\right)v} f_1(s_1 | s_2) \frac{a_2 s_3 e^{-\frac{(\sqrt{2b_2 s_2 - \frac{a_2 s_3}{\sqrt{2}}})^2}{2s_2}}}{2\sqrt{\pi s_2^3}} \times \cdots \times \frac{a_{j+1} e^{-\frac{(\sqrt{2b_{j+1} s_{j+1} - \frac{a_{j+1}}{\sqrt{2}}})^2}{2s_{j+1}}}}{2\sqrt{\pi s_{j+1}^3}} ds_{j+1} \cdots ds_1. \end{aligned}$$

Hence, Equation (2) also holds for $n = j + 1$, which means that this statement holds for any $n \in \mathbb{N}^+$. \square

With the integral representation Equation (2), we can develop an associate simulation scheme for a variety class of RVs by setting different values for the parameters a_k, b_k , and c_k . For instance, if we set $c_k = 0$ for all k , we can develop a general simulation scheme to sample an RV S , where the Laplace transform is of the following form:

$$\mathbb{E} \left[e^{-vS} \right] = a_n \sqrt{\cdots \sqrt{\cdots \sqrt{b_2 + \hat{f}_1(v)} - a_2 \sqrt{b_2} - a_{n-1} \sqrt{b_{n-1}} - a_n \sqrt{b_n}}, \quad (3)$$

when the initial simulation scheme for the RV with Laplace transform $\hat{f}_1(v)$ is known. Let f_1 be the density of RV with Laplace transform $\hat{f}_1(v)$, then, the full simulation structure in general is summarised in Algorithm 1.

⁶The distributional properties of IG distributions have been well documented in Chhikara and Folks (1989).

ALGORITHM 1: General Framework for Simulating S with Laplace Transform Equation (3)

```

(1) if( $n == 1$ ) {
(2)     sample  $S_1$  with distribution  $f_1(s_1)$ 
(3) }
(4) else {
(5)     for  $i \in \{n-1, \dots, 2\}$  {
(6)         sample  $S_n \sim IG(a_n/2\sqrt{b_n}, a_n^2/2)$ 
(7)         set  $a_i = a_k$  and  $b_i = b_k$ 
(8)         sample  $S_i \sim IG(S_{i+1}/2\sqrt{b_i}, a_i^2 S_{i+1}^2/2)$  and sample  $S_1$  with distribution  $f_1(s_1|S_2)$ 
(9)     }
(10) return  $S_1$ 

```

Alternatively, if we set $b_k = c_k = 0$ for all k , this integral representation then leads to an exact simulation scheme for a class of RVs with Laplace transform

$$a_n \sqrt{\cdots \sqrt{\cdots a_2 \sqrt{\hat{f}_1(v)}}} \quad (4)$$

More specifically, if we set $a_k = 1$, for all k and let $\hat{f}_1(v) = v^q$, then, the Laplace transform Equation (1) corresponds to a class of stable distributions with stability index $\frac{q}{2n}$. Indeed, this representation for the stable distributions is of great interest, as it provides us an alternative simulation framework to generate the associated TS class. In the rest of this article, we concentrate on the exact simulation for the TS class with stability index $\frac{q}{2n}$, which is mainly demonstrated in Section 3.

3 TEMPERED STABLE DISTRIBUTIONS AND EXACT SIMULATION

Positive TS distribution is a one-sided tilted stable distribution with stability index $\alpha \in (0, 1)$. It is an *infinitely divisible* distribution that combines both of the stable distribution and Gaussian trends by tilting the *Lévy measure* with an exponential function. More precisely, it has a very flexible structure with three parameters, and the Lévy measure ν is specified by

$$\nu(ds) = e^{-\beta s} \frac{\theta}{s^{\alpha+1}} \mathbf{1}_{\{s>0\}} ds, \quad \alpha \in (0, 1), \quad \beta, \theta \in (0, \infty), \quad (5)$$

where α is the *stability index*, θ is the *intensity parameter*, and β is the *tilting parameter*. In particular, if $\alpha = \frac{1}{2}$, it reduces to a very important distribution, the IG distribution.

In the sequel, for simplicity, this TS distribution uniquely determined by Equation (5) is denoted by $TS(\alpha, \beta, \theta)$, and its probability density function (PDF) is denoted by $f(s; \alpha, \beta, \theta)$. Obviously, if $\beta = 0$, it then returns to a standard positive stable distribution $S(\alpha, \theta)$ with Lévy measure

$$\nu(ds) = \frac{\theta}{s^{\alpha+1}} \mathbf{1}_{\{s>0\}} ds,$$

and density $f(s; \alpha, 0, \theta)$. Proposition 3.1 below illustrates an important and well-known connection between these two distributions.

PROPOSITION 3.1. *The Laplace transform of $TS(\alpha, \beta, \theta)$ is given by*

$$\mathbb{E} \left[e^{-v TS(\alpha, \beta, \theta)} \right] = \exp \left(-\frac{\theta \Gamma(1-\alpha)}{\alpha} \left[(\beta+v)^\alpha - \beta^\alpha \right] \right), \quad v \in \mathbb{R}^+, \quad (6)$$

with the density

$$f(s; \alpha, \beta, \theta) = \exp\left(\frac{\theta\Gamma(1-\alpha)}{\alpha}\beta^\alpha - \beta s\right) \times f(s; \alpha, 0, \theta), \quad (7)$$

where $\Gamma(u)$ is the gamma function, i.e., $\Gamma(u) := \int_0^\infty s^{u-1} e^{-s} ds$.

As the name *stable* implies in stable distribution, stability index α is the most crucial parameter, since this fundamentally determines its distributional property of the so-called *stable laws* (or α -stable): a sum of any two independent stable RVs with the same index α is still a stable RV with index α , and this invariance property does not hold across different α 's (Borak et al. 2005, p. 22). It has been commonly recognised that, the density functions for general specifications on α are hard to be obtained analytically, and this indeed poses the greatest challenge to the further study of their distributional properties and statistical inference. Nevertheless, a different specification for α leads to a different family of stable distributions, and some are particularly attractive in respect of their distinctive distributional properties and potentials for applications. Many of these interesting special families have been extensively investigated in the literature, see Brown and Tukey (1946), Mitra (1981, 1982), Montroll and Bendler (1984), and Penson and Górska (2010); also see surveys in Holt and Crow (1973), Devroye (1986), Zolotarev (1986), Samoradnitsky and Taqqu (1994), and Uchaikin and Zolotarev (1999). For example, the $\frac{1}{2^n}$ -family of stable distributions for $n = -1, 0, 1, \dots$ is particularly prestigious, and the first three members are all famous: they are Gaussian, Cauchy, and Lévy, respectively, and each one has an explicit density function with a tailored algorithm available for exact simulation (Uchaikin and Zolotarev 1999, p. 214–216). However, explicit densities beyond these three members rely on heavy use of the hypergeometric function, which are not ideally suited to our goal for exact simulation.

In the following parts of this section, we first set up our proposed framework of exact simulation for the TS families with stability index that can be expressed in a general dyadic form of

$$\alpha = \frac{q}{2^n} \in (0, 1), \quad q \in \mathbb{N}^+, \quad n \in \{k \in \mathbb{N}^+ \mid 2^k > q\}. \quad (8)$$

We mainly consider the case where q is an odd number as, obviously, an even numerator will reduce to an odd numerator by moving down the degree n . Meanwhile, we focus on some of its important subclasses where further enhanced algorithms (such as *direct* simulation procedure) are available. The associated numerical examples will be presented later in Section 4.

3.1 Tempered Stable Distributions with Stability Index $\alpha = \frac{q}{2^n}$

Based on Theorem 2.1, if $a_k = 1$ and $b_k = c_k = 0$ for any $k = 2, \dots, n$ and $\hat{f}_1(v) = v^q$, then, Equation (1) represents the Laplace transform of a stable distribution with stability index $\frac{q}{2^n}$. Therefore, we can represent the density of a TS distribution with stability index $\frac{q}{2^n} \in (0, 1)$ based on Equations (7) and (2) in Theorem 3.2 as follows.

THEOREM 3.2. For $q, n \in \mathbb{N}^+$ specified by Equation (8) and $p := \min_{1 \leq i \leq n} \{i \in \mathbb{N}^+ : 2^i \geq q\}$, the density of $TS(\frac{q}{2^n}, \beta, \theta)$ has an integral representation of

$$f(s_1) = \begin{cases} \exp(-\beta s_1 + A_p \beta \frac{q}{2^p}) f(s_1; \frac{q}{2^p}, 0, \theta), & n = p, \\ \int_0^\infty \dots \int_0^\infty \frac{\frac{s_2}{\sqrt{2}} e^{-\frac{(\sqrt{2}\beta s_1 - \frac{s_2}{\sqrt{2}})^2}{2s_1}}}{\sqrt{2\pi s_1^3}} \times \dots \times \frac{\frac{s}{\sqrt{2}} e^{-\frac{(\sqrt{2}\beta \frac{1}{2^{n-p}} s_{n-p} - \frac{s}{\sqrt{2}})^2}{2s_{n-p}}}}{\sqrt{2\pi s_{n-p}^3}} f_S(s) e^{-\beta \frac{1}{2^{n-p}} s + A_n \beta \frac{q}{2^n}} ds \dots ds_2, & n > p, \end{cases} \quad (9)$$

where $A_n := \frac{2^n}{q} \theta \Gamma(1 - \frac{q}{2^n})$, and $f_S(s)$ is the density of a stable distribution with stability index $\frac{q}{2^p}$ such that its Laplace transform is $\exp(-A_n v \frac{q}{2^p})$.

When $n = p$, i.e., n is the smallest integer for $\frac{q}{2^n} < 1$, the density is equivalent to Equation (7), and the simulation scheme degenerates to the SSR scheme (with the algorithm in Appendix A). Whereas when $n > p$, from the multiple integral representation Equation (9) for the density function, we can observe that $TS(\frac{q}{2^n}, \beta, \theta)$ is closely related to an IG distribution, which can be directly simulated by Michael et al. (1976).⁷ We therefore develop an alternative simulation framework based on the integral representation for $TS(\frac{q}{2^n}, \beta, \theta)$ in Algorithm 2, namely the *BR scheme*. The recursion first needs to be initialised by generating an RV (named “seed” throughout this article) \tilde{S} via the A/R scheme, the general BR scheme in Algorithm 2 as below to sample $TS(\frac{q}{2^n}, \beta, \theta)$.

ALGORITHM 2: Backward Recursive Scheme for Simulating $TS(\frac{q}{2^n}, \beta, \theta)$

- (1) **set** $\tilde{\theta} := 2^{n-p} \theta \Gamma(1 - \frac{q}{2^n}) / \Gamma(1 - \frac{q}{2^p})$
 - (2) **repeat** {
 - (3) **sample** $\tilde{S} \sim S(q/2^p, \tilde{\theta})$ and $V \sim \mathcal{U}[0, 1]$
 - (4) **if** ($V \leq \exp(-\beta^{1/2^{n-p}} \tilde{S})$) **break**
 - (5) }
 - (6) **set** $\tilde{S}_{n-p+1} = \tilde{S}$
 - (7) **for** $i \in \{n-p, \dots, 2, 1\}$
 - (8) **sample** $\tilde{S}_i \sim IG(\tilde{S}_{i+1}/2\beta^{1/2^i}, \tilde{S}_{i+1}^2/2)$
 - (9) }
 - (10) **return** \tilde{S}_1
-

This alternative scheme turns out to be useful for a large n , in the way that it avoids the low acceptance rate of sampling a TS distribution with a lower stability index by generating a TS distribution with a relative higher stability index and using a recursive procedure to produce the rest. To illustrate the key idea of recursion, we could adopt the SSR scheme to generate the seed, therefore, the corresponding expected complexity of the algorithm based on this SSR scheme is $n - p + 2 \exp(A_p \beta \frac{q}{2^p})$. Since the complexity is exponentially increasing with respect to the parameter β , to accelerate the algorithms, one should replace the SSR algorithm by the DR method (Devroye 2009), or, the fast rejection algorithm (Hofert 2011b), to speed up the simulation of the associated seed.

⁷Note that, the exact simulation for IG distribution designed by Michael et al. (1976) is *direct* and very efficient, as it is developed based on the explicit inverse transformation of multiple roots without the A/R scheme.

3.2 Tempered Stable Distributions with Stability Index $\alpha = \frac{1}{2^n}$

Mathematically elegant schemes of *direct* exact simulation without the A/R mechanism, like the classical CMS method (Chambers et al. 1976), are indeed very rare in the families of TS distributions but extremely desirable. In this section, we present our first discovery of such a unique family that allows *direct* simulation, that is, the family of TS distributions with the stability index of *binary fractions* $\frac{1}{2^n}$, $n = 1, 2, \dots$. In fact, it inherits from the prestigious family of stable distributions with the same style as we mentioned at the beginning of this section. The first member of this $\frac{1}{2^n}$ -family of TS distributions is familiar: $TS(\frac{1}{2}, \beta, \theta)$ follows an IG distribution, since the associated stable distribution $S(\frac{1}{2}, \theta)$ follows a Lévy distribution. However, it is almost unknown for the remaining members.

This family is very unique, which allows us to develop an extremely fast algorithm for exactly sampling any member based on recursively calling the IG generator developed by Michael et al. (1976). It is accurate and very straightforward to implement. Amazingly, it barely depends on the choices of parameters, as the recursion is only driven by the efficient IG generator and no A/R mechanism is involved. This *direct* procedure is summarised in Algorithm 3, and it is derived immediately from a multiple integral representation of its density function in Corollary 3.3.

COROLLARY 3.3. *The density of $TS(\frac{1}{2^n}, \beta, \theta)$ has an integral representation of*

$$f\left(s_1; \frac{1}{2^n}, \beta, \theta\right) = \int_0^\infty \cdots \int_0^\infty \frac{\frac{s_2}{\sqrt{2}}}{\sqrt{2\pi s_1^3}} e^{-\frac{\left(\sqrt{2\beta} s_1 - \frac{s_2}{\sqrt{2}}\right)^2}{2s_1}} \times \cdots \times \frac{\frac{A_n}{\sqrt{2}}}{\sqrt{2\pi s_n^3}} e^{-\frac{\left(\sqrt{2\beta} \frac{1}{2^n} s_n - \frac{A_n}{\sqrt{2}}\right)^2}{2s_n}} ds_n \cdots ds_2, \quad (10)$$

where $A_n = 2^n \theta \Gamma(1 - \frac{1}{2^n})$.

Proof for Corollary 3.3 immediately follows Theorem 2.1 by setting $\hat{f}_1(v) = \sqrt{v}$. Hence, based on Equation (10), we can directly generate a $TS(\frac{1}{2^n}, \beta, \theta)$ RV for $n > 1$ via the following Algorithm 3.

ALGORITHM 3: Direct Scheme for Simulating $TS(\frac{1}{2^n}, \beta, \theta)$

- (1) **sample** $\tilde{S}_{n+1} \sim IG\left(2^n \theta \Gamma(1 - \frac{1}{2^n})/2\beta^{1/2^n}, [2^n \theta \Gamma(1 - 1/2^n)]^2/2\right)$
 - (2) **for** $i \in \{n, \dots, 2, 1\}$
 - (3) **sample** $\tilde{S}_i \sim IG\left(\tilde{S}_{i+1}/2\beta^{1/2^{i-1}}, \tilde{S}_{i+1}^2/2\right)$
 - (4) }
 - (5) **return** \tilde{S}_1
-

To illustrate how Algorithm 3 works, let us consider the simplest case when $n = 2$ as an example: Since $TS(\frac{1}{4}, \beta, \theta)$ is equal in distribution to \tilde{S}_1 , it can be simulated by first generating an IG RV of

$$\tilde{S}_2 \sim IG\left(\frac{2\theta\Gamma\left(\frac{3}{4}\right)}{\beta^{\frac{1}{4}}}, 8\theta^2\Gamma^2\left(\frac{3}{4}\right)\right);$$

then, conditional on one realisation of \tilde{S}_2 , we can generate $TS(\frac{1}{4}, \beta, \theta)$ i.e., \tilde{S}_1 via another IG RV of

$$\tilde{S}_1 \mid \tilde{S}_2 \sim IG\left(\frac{\tilde{S}_2}{2\sqrt{\beta}}, \frac{\tilde{S}_2^2}{2}\right).$$

3.3 Tempered Stable Distributions with Stability Index $\alpha = \frac{3}{2n}$

To sample TS RVs beyond the $\frac{1}{2n}$ -family, we find that it is hard to obtain the associated *direct* algorithms and eventually the A/R scheme is unavoidable. Similar to the stable distribution, there is no explicit formula for the density of TS distribution in general. However, for some special choices of q , we find an alternative way to generate the stable distribution $S(\frac{q}{2p}, \tilde{\theta})$ of the seed without using *Zolotarev's integral representation*. In this section, we take a further inside of the density of stable distribution of $S(\frac{3}{4}, \theta)$ based on some very interesting distributional results we discovered in Lemma 3.4 for a special family of RVs with Laplace transform Equation (11). Then, we design an enhanced algorithm to generate the $\frac{3}{2n}$ -TS family via algorithm for $n = 2$ and algorithm for $n > 2$, respectively.

Let us first introduce a class of RVs with Laplace transform Equation (11), which is a simplified version of Equation (1). This class of RVs has a closed-form distribution function, which leads to a direct simulation scheme. Simply setting $\gamma = 0$, Equation (11) becomes the Laplace transform of an $S(\frac{3}{4}, \theta)$ RV. Details of the distribution property and simulation scheme for this special family are presented in Lemma 3.4 and Algorithm 4.

LEMMA 3.4. *If the RV S has the Laplace transform*

$$\mathbb{E} \left[e^{-vS} \right] = \exp \left(-\zeta \sqrt{v^{\frac{3}{2}} + \gamma v} \right), \quad \zeta, \gamma \in \mathbb{R}^+, \quad (11)$$

then, the corresponding CDF of S can be represented by

$$F_S(s) = \int_0^\infty \frac{1}{\pi(w+1)\sqrt{w}} \exp \left(-\frac{\zeta^4(w+1)^3}{64w} \frac{1}{s^3} - \frac{\gamma \zeta^2(w+1)}{4} \frac{1}{s} \right) dw. \quad (12)$$

PROOF. Let $f_S(s)$ and $F_S(s)$ denote the PDF and the cumulative distribution function (CDF) of S , respectively. According to Equation (11), the Laplace transform of $F_S(s)$ can be derived:

$$\begin{aligned} \int_0^\infty e^{-vs} F_S(s) ds &= \int_0^\infty \int_u^\infty e^{-vs} f_S(u) ds du \\ &= \frac{1}{v} \int_0^\infty e^{-vu} f_S(u) du \\ &= \frac{1}{v} \exp \left(-\zeta \sqrt{v^{\frac{3}{2}} + \gamma v} \right), \quad v \in \mathbb{R}^+. \end{aligned} \quad (13)$$

Note that, by change of variable $u = vx$, Equation (13) can be rewritten as

$$\begin{aligned} \frac{1}{v} \exp \left(-\zeta \sqrt{v^{\frac{3}{2}} + \gamma v} \right) &= \frac{1}{v} \int_0^\infty e^{-(\sqrt{v}-\gamma)vx} \frac{\zeta}{2\sqrt{\pi x^3}} \exp \left(-\frac{\zeta^2}{2x} \right) dx \\ &= \frac{1}{\sqrt{v}} \int_0^\infty e^{-\sqrt{v}u} \frac{\zeta}{2\sqrt{\pi u^3}} \exp \left(-\frac{\zeta^2 v}{4u} - \gamma u \right) du. \end{aligned}$$

From the tables of Laplace transforms listed in Bateman (1954, p. 246), we recognise that

$$\frac{1}{\sqrt{v}} e^{-\sqrt{v}u} = \int_0^{\infty} e^{-vy} \frac{1}{\sqrt{\pi y}} \exp\left(-\frac{u^2}{4y}\right) dy.$$

Finally, Equation (13) has an integral representation of

$$\begin{aligned} \frac{1}{v} \exp\left(-\zeta\sqrt{v^{\frac{3}{2}} + \gamma v}\right) &= \int_0^{\infty} \int_0^{\infty} e^{-vy} \frac{1}{\sqrt{\pi y}} \exp\left(-\frac{u^2}{4y}\right) \frac{\zeta}{2\sqrt{\pi u^3}} \exp\left(-\frac{\zeta^2 v}{4u} - \gamma u\right) dudy \\ &= \int_0^{\infty} \int_0^{\infty} e^{-vy} e^{vz} \frac{1}{\pi\sqrt{yz}} \exp\left(-\frac{\zeta^4}{64yz^2} - \frac{\gamma\zeta^2}{4z}\right) dzdy. \end{aligned} \quad (14)$$

Applying transformations $s = y + z$ and $w = \frac{y}{z}$ to Equation (14), we have $z = \frac{s}{w+1}$, $y = \frac{sw}{w+1}$ with the Jacobian of the transformation

$$\left| \begin{array}{cc} \frac{1}{w+1} & \frac{w}{w+1} \\ -\frac{s}{(w+1)^2} & \frac{s}{(w+1)^2} \end{array} \right| = \frac{s}{(w+1)^2}.$$

Hence, Equation (13) can be finally rewritten as

$$\frac{1}{v} \exp\left(-\zeta\sqrt{v^{\frac{3}{2}} + \gamma v}\right) = \int_0^{\infty} \int_0^{\infty} e^{-vs} \frac{1}{\pi(1+w)\sqrt{w}} \exp\left(-\frac{\zeta^4(1+w)^3}{64w} \frac{1}{s^3} - \frac{\gamma\zeta^2(1+w)}{4} \frac{1}{s}\right) ds,$$

which corresponds to the CDF of Equation (12). \square

According to the closed-form CDF Equation (12) for S with Laplace transform Equation (11), we can directly generate S using an explicit inverse transformation. The simulation scheme for sampling S is provided in Algorithm 4 as follows.

ALGORITHM 4: Exact Scheme for Simulating S with Laplace Transform Equation (11)

- (1) **sample** $U^{(1)} \sim \mathcal{U}[0, 1]$, and $U^{(2)} \sim \mathcal{U}[0, 1]$
- (2) **set** $W = \tan^2\left(\frac{\pi}{2}U^{(1)}\right)$, $A = \frac{\zeta^4(1+W)^3}{64W}$, $B = \frac{\gamma\zeta^2(1+W)}{4}$, $C = -\ln(U_2)$, and

$$D = 3\sqrt{3}\sqrt{27A^2C^4 + 4AB^3C^2 + 27AC^2 + 2B^3}$$

- (3) **return**

$$\frac{1}{3C} \left(D^{\frac{1}{3}}/2^{\frac{1}{3}} + 2^{\frac{1}{3}}B^2/D^{\frac{1}{3}} + B \right) \quad (15)$$

PROOF. We observe that the CDF of S in Equation (12) of Lemma 3.4 can be represented by

$$F_S(s) = \int_0^{\infty} f_W(w) F_{S|W}(s|w) dw,$$

where W is a well-defined RV with the density

$$f_W(w) = \frac{1}{\pi\sqrt{w(w+1)}}.$$

The corresponding CDF is

$$F_W(w) = \int_0^w \frac{1}{\pi\sqrt{x}(1+x)} dx = \frac{2}{\pi} \tan^{-1}(\sqrt{w}),$$

which allows W to be directly sampled via an explicit inverse transform by setting

$$W \stackrel{\mathcal{D}}{=} \tan^2\left(\frac{\pi}{2}U^{(1)}\right), \quad U^{(1)} \sim \mathcal{U}[0, 1].$$

Conditional on W , the CDF of S is therefore given by

$$F_{S|W}(s|w) = \exp\left(-\frac{\zeta^4(1+w)^3}{64w} \frac{1}{s^3} - \frac{\gamma\zeta^2(1+w)}{4} \frac{1}{s}\right).$$

To sample S given W , we then need to solve the following cubic equation:

$$-\ln(U_2) \times S^3 - \frac{\gamma\zeta^2(1+W)}{4} \times S^2 - \frac{\zeta^4(1+W)^3}{64W} = 0, \quad U^{(2)} \sim \mathcal{U}[0, 1]. \quad (16)$$

The solution for this cubic equation immediately follows Equation (15) using the parameter setting in Algorithm 4 for the coefficients in Equation (16). \square

For a stable RV $S(\frac{3}{4}, \theta)$, its Laplace transform satisfies Equation (11) with $\gamma = 0$. The corresponding distribution function for $S(\frac{3}{4}, \theta)$ satisfies Equation (17).

COROLLARY 3.5. *The CDF of stable distribution $S(\frac{3}{4}, \theta)$ for any $\theta \in \mathbb{R}^+$ can be represented by*

$$F_{S(\frac{3}{4}, \theta)}(s) = \int_0^\infty \frac{1}{\pi(w+1)\sqrt{w}} \exp\left(-\frac{\kappa^4(w+1)^3}{64s^3w}\right) dw, \quad (17)$$

and the simulation scheme for the stable RV $S(\frac{3}{4}, \theta)$ directly follows Algorithm 4 by setting $B = 0$.

PROOF. The Laplace transform of $S(\frac{3}{4}, \theta)$ is given by

$$\mathbb{E}\left[e^{-vS(\frac{3}{4}, \theta)}\right] = e^{-\kappa v^{\frac{3}{4}}}, \quad \kappa = \frac{4}{3}\theta\Gamma\left(\frac{1}{4}\right).$$

Based on Lemma 3.4, setting $\gamma = 0$ and $\zeta = \kappa$ in Equation (12), the CDF of $S(\frac{3}{4}, \theta)$ follows Equation (17). \square

To generate the associated TS distribution $TS(\frac{3}{4}, \beta, \theta)$, instead of using the A/R based on Zolotarev's integral representation, we redesign an alternative A/R scheme using a gamma distributed envelop based on the distribution function we obtained for the stable distribution $S(\frac{3}{4}, \theta)$. The detail is given in Algorithm 5.

ALGORITHM 5: A/R Scheme for Simulating $TS(\frac{3}{4}, \beta, \theta)$

- (1) **set** $\kappa = 4\theta\beta^{\frac{3}{4}}\Gamma\left(\frac{1}{4}\right)/3$
 - (2) **repeat** {
 - (3) **sample** $W = \tan^2\left(\frac{\pi}{2}U\right)$ with $U \sim \mathcal{U}[0, 1]$
 - (4) **sample** $E \sim \Gamma(m, 1)$ and $V \sim \mathcal{U}[0, 1]$
 - (5) **if** $(V \leq 3^{m+1}6^{\frac{m}{3}}\kappa^{\frac{4(m+3)}{3}}e^{\frac{m+3}{3}}(1+W)^3e^{-\frac{\kappa^4(1+W)^3}{64WE^3}} / (2^{3(m+2)}(m+3)^{\frac{m+3}{3}}WE^{m+3}))$ **break**
 - (6) }
 - (7) **return** E/β
-

PROOF. Let $X \sim TS(\frac{3}{4}, \beta, \theta)$ and set $S = \beta X$. According to Proposition 3.1, the Laplace transform of S is given by

$$\begin{aligned} \mathbb{E} \left[e^{-vS} \right] &= \mathbb{E} \left[e^{-v\beta X} \right] \\ &= \exp \left(-\frac{4\theta\Gamma(\frac{1}{4})}{3} \left[(\beta + v\beta)^{\frac{3}{4}} - \beta^{\frac{3}{4}} \right] \right) \\ &= \exp \left(-\frac{4\theta\beta^{\frac{3}{4}}\Gamma(\frac{1}{4})}{3} \left[(1+v)^{\frac{3}{4}} - 1 \right] \right) \\ &= \exp \left(-\kappa \left[(1+v)^{\frac{3}{4}} - 1 \right] \right), \end{aligned}$$

where $\kappa = \frac{4}{3}\theta\beta^{\frac{3}{4}}\Gamma(\frac{1}{4})$. Using Equations (7) and (17), the density of S can be written as

$$f_S(s) = \int_0^\infty \frac{1}{\pi(w+1)\sqrt{w}} \frac{3\kappa^4(1+w)^3}{64ws^4} \exp \left(-\frac{\kappa^4(1+w)^3}{64ws^3} \right) e^{-s+\kappa} dw. \quad (18)$$

The joint density of S and W is

$$f(s, w) = \frac{1}{\pi(w+1)\sqrt{w}} \frac{3\kappa^4(1+w)^3}{64ws^4} \exp \left(-\frac{\kappa^4(1+w)^3}{64ws^3} \right) e^{-s+\kappa}, \quad w, s > 0.$$

To generate S , we consider a Gamma distributed envelop with rate parameter $m > 0$ and scale parameter 1 for S , and this envelop appears to be the most natural choice to be adopted according to its density Equation (18). The joint density we have is

$$g(s, w) = \frac{1}{\pi(w+1)\sqrt{w}} \frac{1}{\Gamma(m)} s^{m-1} e^{-s}, \quad w, s > 0.$$

We have

$$\begin{aligned} \frac{f(s, w)}{g(s, w)} &= \frac{3\kappa^4(w+1)^3\Gamma(m)e^\kappa}{64w} \frac{1}{s^{3+m}} \exp \left(-\frac{\kappa^4(1+w)^3}{64ws^3} \right) \\ &\leq \frac{\Gamma(m)e^\kappa e^{-\frac{m+3}{3}} \left(\frac{64}{3\kappa^4} \right)^{\frac{m}{3}} (m+3)^{\frac{m+3}{3}} w^{\frac{m}{3}}}{(w+1)^m} \\ &\leq \Gamma(m)e^\kappa e^{-\frac{m+3}{3}} \left(\frac{8}{3} \right)^m \left(\frac{1}{6\kappa^4} \right)^{\frac{m}{3}} (m+3)^{\frac{m+3}{3}} = C(m, \kappa), \end{aligned}$$

where $C(m, \kappa)$ can be minimised over m . It is equivalent to find the m that minimises

$$\log C(m, \kappa) = \frac{1}{3} [\log \Gamma(m) + 3\kappa - (m+3) + 8m \log(2) - 4m \log(3) - 4m \log(\kappa) + (m+3) \log(m+3)].$$

The optimal value m^* satisfies the following equation:

$$\frac{1}{4} \left[\log(m^* + 3) + 3\psi^{(0)}(m^*) \right] = \log \left(\frac{3\kappa}{4} \right), \quad \text{for } \psi^{(0)}(m) = \frac{d\Gamma(m)}{dm},$$

and we have

$$m^* = \frac{3\kappa}{4} - \frac{3}{8} + \frac{143}{128} \frac{1}{m^*} - \frac{1857}{1024} \frac{1}{m^{*2}} + O \left(\frac{1}{m^{*3}} \right) \approx \frac{3\kappa}{4} + c.$$

When we numerically fit this linear setting, the optimal value c obtained is -0.14 . Hence, given κ , the optimal rate m^* for the gamma distributed envelop is chosen by setting

$$m^* = -0.14 + \frac{3}{4}\kappa.$$




Fig. 1. Acceptance rates of Algorithm 5 and the SSR scheme for sampling $TS(\frac{3}{4}, \beta, \theta)$ based on the parameter setting $\theta = 0.7$ and $\beta \in (0, 10)$.

To generate S , we first sample W using the explicit inverse transformation, and then generate the gamma distribution envelop $E \sim \Gamma(m^*, 1)$. The A/R decision follows

$$V \leq \frac{f(E, W)}{C(m^*, \kappa)g(E, W)} = \frac{3^{m^*+1} 6^{\frac{m^*}{3}} \kappa^{\frac{4(m^*+3)}{3}} e^{\frac{m^*+3}{3}} (1+W)^3 e^{-\frac{\kappa^4(1+W)^3}{64WE^3}}}{2^{3(m^*+2)} (m^*+3)^{\frac{m^*+3}{3}} W E^{m^*+3}},$$

and the corresponding acceptance rate is $\frac{1}{C(m^*, \kappa)}$. Finally, we obtain $TS(\frac{3}{4}, \beta, \theta)$ by setting $X = \frac{S}{\beta}$. \square

Comparing with the acceptance rate $\exp(-\frac{4}{3}\theta\Gamma(\frac{1}{4})\beta^{\frac{3}{4}})$ using the SSR scheme to generate $TS(\frac{3}{4}, \beta, \theta)$, Algorithm 5 leads to a higher acceptance rate, in particular for a large value of β . The difference of the acceptance rates to generate one $TS(\frac{3}{4}, \beta, \theta)$ RV for these two schemes with respect to β is plotted in Figure 1. The SSR scheme is only desirable when the tilting parameter is small, whereas our Algorithm 5 is more competitive when the tilting parameter is large. Hence, instead of using the SSR scheme to generate the seeds for the $TS(\frac{3}{2^n}, \beta, \theta)$ family, we adopt Algorithm 5 to sample seeds. The enhanced algorithm for $TS(\frac{3}{2^n}, \beta, \theta)$ is provided in Algorithm 6.

PROOF. Letting $X \sim TS(\frac{3}{2^n}, \beta, \theta)$ and $S = \beta X$, we have

$$\begin{aligned} \mathbb{E} [e^{-vS}] &= \mathbb{E} [e^{-v\beta X}] \\ &= \exp\left(-\frac{2^n \theta \Gamma\left(1 - \frac{3}{2^n}\right)}{3} \left[(\beta + \beta v)^{\frac{3}{2^n}} - \beta^{\frac{3}{2^n}}\right]\right) \\ &= \exp\left(-\frac{2^n \theta \beta^{\frac{3}{2^n}} \Gamma\left(1 - \frac{3}{2^n}\right)}{3} \left[(1 + v)^{\frac{3}{2^n}} - 1\right]\right). \end{aligned}$$

ALGORITHM 6: Backward Recursive Scheme for Simulating $TS(\frac{3}{2^n}, \beta, \theta)$

```

(1) set  $\kappa = 2^n \theta \beta^{\frac{3}{2^n}} \Gamma(1 - 3/2^n) / 3$ 
(2) repeat {
(3)   sample  $W = \tan^2(\frac{\pi}{2}U)$  with  $U \sim \mathcal{U}[0, 1]$ 
(4)   sample  $E \sim \Gamma(m, 1)$  and  $V \sim \mathcal{U}[0, 1]$ 
(5)   if  $(V \leq 3^{m+1} 6^{\frac{m}{3}} \kappa^{\frac{4(m+3)}{3}} e^{\frac{m+3}{3}} (1+W)^3 e^{-\frac{\kappa^4(1+W)^3}{64WE^3}} / (2^{3(m+2)}(m+3)^{\frac{m+3}{3}} WE^{m+3}))$  break
(6) }
(7) set  $\tilde{S}_{n-1} = E$ 
(8) for  $i \in \{n-2, \dots, 2, 1\}$ 
(9)   sample  $\tilde{S}_i \sim IG(\tilde{S}_{i+1}/2, \tilde{S}_{i+1}^2/2)$ 
(10) }
(11) return  $\tilde{S}_1/\beta$ 

```

Based on Theorem 3.2 and Equation (18), the density of $S \sim TS(\frac{3}{2^n}, 1, \theta \beta^{\frac{3}{2^n}})$ has the integral representation

$$\begin{aligned}
f\left(s_1; \frac{3}{2^n}, 1, \theta \beta^{\frac{3}{2^n}}\right) &= \int_0^\infty \cdots \int_0^\infty \frac{\frac{s_2}{\sqrt{2}}}{\sqrt{2\pi s_1^3}} e^{-\frac{(\sqrt{2}s_1 - \frac{s_2}{\sqrt{2}})^2}{2s_1}} \times \cdots \times \frac{\frac{s}{\sqrt{2}}}{\sqrt{2\pi s_{n-2}^3}} e^{-\frac{(\sqrt{2}s_{n-2} - \frac{s}{\sqrt{2}})^2}{2s_{n-2}}} \\
&\quad \times \frac{3\kappa^4(w+1)^2}{64\pi w^{\frac{3}{2}} s^4} \exp\left(-\frac{\kappa^4(w+1)^3}{64s^3 w} - s + \kappa\right) dw \cdots ds_2,
\end{aligned}$$

where

$$\kappa = \frac{2^n}{3} \theta \beta^{\frac{3}{2^n}} \Gamma\left(1 - \frac{3}{2^n}\right). \quad (19)$$

The seed, denoted \tilde{S} , follows a distribution of $TS(\frac{3}{4}, 1, \theta \beta^{\frac{3}{2^n}})$, and the density satisfies Equation (18) with the new κ defined in Equation (19). We directly follow Algorithm 5 to generate seeds and follow the general BR scheme to generate X . \square

The total number of RVs needed to generate one $TS(\frac{3}{2^n}, \beta, \theta)$ for our BR scheme would be $n - 2 + 3C(m^*, \kappa)$, which is the sum of the expected number of iterations of the A/R algorithm required to generate a seed and the number of RVs generated via the recursion. The total number of RVs needed for the classical SSR scheme is $2 \exp(\frac{2^n \theta}{3} \Gamma(1 - \frac{3}{2^n}) \beta^{\frac{3}{2^n}})$, so our BR scheme shall have a better performance in general. In Figure 2, we compare the logarithm of the expected number of iterations for these two algorithms. The computation cost for the BR scheme is less than the SSR scheme, especially, when the tilting parameter β becomes larger. In general, different from the SSR scheme, the expected complexity of the BR scheme does not increase exponentially with respect to the β .

4 NUMERICAL EXAMPLES

In this section, we provide numerical examples for these three families of TS distributions with stability index $\alpha = \frac{1}{2^n}, \frac{3}{2^n}, \frac{q}{2^n}$, respectively. The simulation experiments are all conducted on a normal laptop with the Intel Core i7-6500U CPU@2.50GHz processor, 8GB RAM, Windows 10 Home, and 64-bit Operating System. The algorithms are coded and performed in MatLab (R2016b), and the computing time is measured by the *elapsed CPU time* in seconds. Numerical validation and




Fig. 2. Logarithm of the expected number of random variables needed for our *backward recursive* (BR) scheme (via Algorithm 5) versus the *simple stable rejection* (SSR) via Algorithm 7 based on the parameter setting $\theta = 0.7$, $n = 2, 3, 4, 5$, respectively, for the tilting parameter $\beta \in (0, 10)$.

tests for our simulation algorithms are based on the PDF and CDF of $TS(\alpha, \beta, \theta)$, which can be calculated by inverting the Laplace transform Equation (6) numerically.⁸ In particular, we assess the goodness-of-fit by comparing the CDFs and PDFs obtained by our BR scheme and by the numerical inverse of Laplace transform. The associated plots of CDFs, PDFs and histograms under different parameter settings are illustrated in Figures 3, 4, and 5, respectively. Overall, we can observe that each of these algorithms can achieve a very high level of accuracy, and the simulated CDF and PDF are fitted well to the associated numerical inversion.

We carry out a comparison of CPU time for our BR scheme against the SSR scheme and the DR scheme for three families of TS distributions with stability index $\alpha = \frac{1}{2^n}, \frac{3}{2^n}, \frac{q}{2^n}$. The associated numerical results in detail is reported in Table 1. Comparing with the SSR scheme and DR scheme, our direct scheme of Algorithm 3 for the first family of $\alpha = \frac{1}{2^n}$ performs extraordinarily fast, since no A/R procedure is involved as clearly demonstrated in Section 3.2. For example, it is nearly 1,000 times faster than the SSR scheme and 10 times faster than the DR scheme for simulating 100,000 replications based on parameter setting $\alpha = 1/32, \theta = 0.5, \beta = 1$. The out-performance of our algorithm would even become much more substantial when θ or β increases or n increases. More remarkably, the speed of simulation is barely effected by the changes of parameters θ and β , and the CPU time only slightly increases when n increases. The associated numerical results in

⁸A variety of methods are available for numerically inverting Laplace transforms with high accuracy, such as Gaver (1966), Stehfest (1970), Abate and Whitt (1992, 1995, 2006). Here, we use the Euler scheme in Abate and Whitt (2006).




Fig. 3. Comparison of the empirical CDF/PDF for our backward recursive (BR) scheme (via Algorithm 3) with the CDF/PDF obtained via numerical inverse under the parameter setting $\alpha = 1/2^2, 1/2^5, \theta = 0.5, \beta = 3.0$, respectively.

Table 1. Comparison of CPU Time for Backward Recursive (BR), Simple Stable Rejection (SSR), Double Rejection (DR) for $TS(\frac{q}{2^n}, \theta, \beta)$ Based on Parameter Setting $q = 1, 3, 23, 27, 35, n = 2, 3, 5, 6, 7, \theta = 0.50, 0.70, \beta = 1.0, 2.0$

α	$1/2^2$	$1/2^3$	$1/2^5$	$3/2^2$	$3/2^3$	$3/2^5$	$23/2^6$	$27/2^6$	$35/2^7$
	$\theta = 0.50$				$\beta = 1.0$				
BR	1.58	2.20	2.34	1.25	7.62	7.50	4.45	4.36	8.85
SSR	7.45	47.25	2152.53	7.53	6.52	178.39	4.97	4.47	6.15
DR	25.31	23.85	24.89	27.94	23.67	23.48	26.68	25.34	31.52
	$\theta = 0.70$				$\beta = 2.0$				
BR	1.42	2.14	2.87	1.29	8.21	8.29	30.78	45.45	32.53
SSR	54.41	581.38	3272.53	257.76	32.28	3681.59	35.25	39.65	28.93
DR	41.23	37.72	32.89	34.24	30.78	31.45	31.12	32.65	35.12

Each value in the tables is produced from 100,000 replications.

detail for this sensitivity analysis are reported in Table 2. Our Algorithm 6 is also very fast and can easily beat the SSR scheme, as the generator for the stable law of seeds is enhanced by our own design of Algorithm 5. For example, it is about 400 times faster than the SSR scheme for exactly simulating 100,000 replications with parameter setting of $\alpha = 3/32, \theta = 0.7, \beta = 2.0$. Moreover, it is about four times faster than the DR scheme. However, if we generate the seed using the SSR




Fig. 4. Comparison of the empirical CDF/PDF for our backward recursive (BR) scheme (via Algorithms 5 and 6) with the CDF/PDF obtained via numerical inverse under the parameter setting $\alpha = 3/2^2, 3/2^5, \theta = 0.7, \beta = 2.0$, respectively.

scheme for our Algorithm 2 in general for the third family of $\alpha = \frac{q}{2^n}$, then, the performance is at a similar level as the SSR scheme. In general, increasing n would increase the computing time, as more IG variables are needed but the computing time mainly depends on the time of generating the seed. Hence, when the acceptance rate of SSR becomes smaller for a larger tilting parameter β , we can replace the simulation scheme of the seed by DR scheme to improve the simulation speed.

5 CONCLUDING REMARKS

In this article, we provide a new framework for the exact simulation of a class of TS and related distributions based on the multiple integral representation. The principle of our approach, which is based on the backward recursion, is clearly distinguishing from the existing algorithms in the literature. It works pretty efficiently for some subclasses of TS distributions, and is also applicable to some other classes beyond the TS family. This scheme could lead to many prospective applications in practice with improved numerical efficiency: It could be used as the basis to exactly generate TS-constructed RVs. For instance, it is well known that, one two-sided TS RV on the real line \mathbb{R} can be decomposed as the difference of two independent one-sided TS RVs on the positive half line \mathbb{R}^+ (Küchler and Tappe 2013, p. 4262). Moreover, it can also be used to generate paths of various TS-driven stochastic processes, such as *Lévy TS subordinators*; see an abundance of their applications for modelling the stochastic volatility with the associated econometric methods in Barndorff-Nielsen and Shephard (2001b, 2002, 2003), Barndorff-Nielsen et al. (2002), Gander and Stephens (2007a, 2007b), Andrieu et al. (2010), and Todorov et al. (2015). These RVs and stochastic processes could further lead to numerous applications in many other fields. In particular, it would




Fig. 5. Comparison of the empirical CDF/PDF for our backward recursive (BR) scheme (via Algorithm 2) with the CDF/PDF obtained via numerical inverse under the parameter setting $\alpha = 13/2^5, 23/2^6, \theta = 0.25, \beta = 0.5$, respectively.

Table 2. Sensitivity Analysis of the CPU Time Against the Varying Parameters $(\theta, \beta) \in (0, 1] \times (0, 1]$ for Our *Backward Recursive* (BR) Scheme (via Algorithm 3) for $\alpha = 1/2^n, n = 2, 3, 4, 5$, Respectively

$\theta \backslash \beta$	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
$\alpha = 1/4$										
0.2	1.34	1.17	1.23	1.28	1.42	1.19	1.30	1.34	1.30	1.39
0.6	1.16	1.20	1.22	1.14	1.22	1.13	1.13	1.22	1.16	1.30
1.0	1.11	1.14	1.17	1.19	1.13	1.13	1.22	1.14	1.16	1.25
$\alpha = 1/8$										
0.2	2.52	2.19	2.17	2.59	2.33	2.19	2.22	2.30	2.14	2.20
0.6	2.45	2.20	2.19	2.42	2.16	2.13	2.16	2.19	2.25	2.22
1.0	2.50	2.19	2.17	2.41	2.23	2.16	2.25	2.11	2.09	2.11
$\alpha = 1/16$										
0.2	2.91	3.08	2.55	2.47	2.92	2.92	2.58	2.44	2.84	2.48
0.6	2.91	2.91	2.45	2.52	2.91	3.19	2.47	2.52	2.89	2.53
1.0	2.91	2.88	2.42	2.52	2.88	2.88	2.48	2.50	2.89	2.64
$\alpha = 1/32$										
0.2	2.88	2.81	2.88	3.33	2.94	2.80	2.81	2.81	3.38	2.91
0.6	2.88	3.05	2.92	3.45	2.94	2.81	2.84	2.80	3.27	2.89
1.0	2.88	2.80	2.81	3.28	2.84	2.73	2.80	2.73	3.31	2.84

Each value in the tables is produced from 100,000 replications.

be extremely useful for numerical validating and testing the newly developed statistical inference or econometric methods for the TS-based models, where the parameters are set up for Monte Carlo studies. Our Algorithm 3 for the $\frac{1}{2^n}$ -family is strongly recommended to be adopted for generating the required data, and a variety of choices for the parameters n , β and θ would provide enough flexibility for the purpose of numerical testing and validation. Since it is a bit of an art to find appropriate envelopes in the development of A/R schemes, the further enhancement for these A/R schemes adopted in this article could be a meaningful topic for future research.

APPENDIX

A SIMPLE STABLE REJECTION (SSR) SCHEME

The SSR scheme introduced by Brix (1999) is illustrated in Algorithm 7 as below.

ALGORITHM 7: Simple Stable Rejection (SSR) Scheme for $TS(\alpha, \beta, \theta)$

- (1) **repeat**{
 - (2) **sample** $U \sim \mathcal{U}[0, \pi]$, $E_s \sim \text{Exp}(1)$ and $V \sim \mathcal{U}[0, 1]$
 - (3) **set** $S = (-\theta\Gamma(-\alpha))^{-\frac{1}{\alpha}} \sin(\alpha U + \pi\alpha/2) \cos(U)^{-\frac{1}{\alpha}} \cos((1-\alpha)U - \pi\alpha/2)^{\frac{1-\alpha}{\alpha}} E_s^{-\frac{1-\alpha}{\alpha}}$
 - (4) **if** ($V \leq \exp(-\beta S)$) **break**
 - (5) }
 - (6) **return** S
-

ACKNOWLEDGMENTS

The authors would like to thank the associate editor and reviewers for many constructive comments that greatly improved the presentation of our results.

REFERENCES

- J. Abate and W. Whitt. 1992. The Fourier-series method for inverting transforms of probability distributions. *Queueing Systems* 10, 1 (1992), 5–87.
- J. Abate and W. Whitt. 1995. Numerical inversion of Laplace transforms of probability distributions. *ORSA Journal on Computing* 7, 1 (1995), 36–43.
- J. Abate and W. Whitt. 2006. A unified framework for numerically inverting Laplace transforms. *INFORMS Journal on Computing* 18, 4 (2006), 408–421.
- C. Andrieu, A. Doucet, and R. Holenstein. 2010. Particle Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 72, 3 (2010), 269–342.
- B. Baeumer and M. M. Meerschaert. 2010. Tempered stable Lévy motion and transient super-diffusion. *Journal of Computational and Applied Mathematics* 233, 10 (2010), 2438–2448.
- O. E. Barndorff-Nielsen, E. Nicolato, N. Shephard, et al. 2002. Some recent developments in stochastic volatility modelling. *Quantitative Finance* 2, 1 (2002), 11–23.
- O. E. Barndorff-Nielsen and N. Shephard. 2001a. Non-Gaussian Ornstein–Uhlenbeck-based models and some of their uses in financial economics. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 63, 2 (2001a), 167–241.
- O. E. Barndorff-Nielsen and N. Shephard. 2001b. Normal modified stable processes. *Theory of Probability and Mathematical Statistics*, pages 1–19.
- O. E. Barndorff-Nielsen and N. Shephard. 2002. Econometric analysis of realized volatility and its use in estimating stochastic volatility models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 64, 2 (2002), 253–280.
- O. E. Barndorff-Nielsen and N. Shephard. 2003. Integrated OU processes and non-Gaussian OU-based stochastic volatility models. *Scandinavian Journal of Statistics* 30, 2 (2003), 277–295.
- O. E. Barndorff-Nielsen, T. Mikosch, and S. I. Resnick. 2012. *Lévy Processes: Theory and Applications*. Springer Science & Business Media.
- H. Bateman. 1954. *Tables of Integral Transforms*, Vols. I & II McGraw-Hill.
- L. Bondesson. 1982. On simulation from infinitely divisible distributions. *Advances in Applied Probability* 14, 4 (1982), 855–869.

- S. Borak, W. Härdle, and R. Weron. 2005. Stable distributions. In *Statistical Tools for Finance and Insurance*, P. Cizek, W. K. Härdle, and R. Weron (Eds.). Springer, Berlin, Heidelberg, pages 21–44.
- A. Brix. 1999. Generalized gamma measures and shot-noise Cox processes. *Advances in Applied Probability* 31, 4 (1999), 929–953.
- G. W. Brown and J. W. Tukey. 1946. Some distributions of sample means. *The Annals of Mathematical Statistics* 17, 1 (1946), 1–12.
- P. Carr and L. Wu. 2003. The finite moment log stable process and option pricing. *The Journal of Finance* 58, 2 (2003), 753–777.
- J. M. Chambers, C. L. Mallows, and B. W. Stuck. 1976. A method for simulating stable random variables. *Journal of the American Statistical Association* 71, 354 (1976), 340–344.
- R. Chhikara and L. Folks. 1989. *The Inverse Gaussian Distribution: Theory, Methodology, and Applications*. Marcel Dekker, New York.
- R. Cont and P. Tankov. 2004. *Financial Modelling with Jump Processes*. CRC Press.
- L. Devroye. 1986. *Non-Uniform Random Variate Generation*. Springer-Verlag.
- L. Devroye. 2009. Random variate generation for exponentially and polynomially tilted stable distributions. *ACM Transactions on Modeling and Computer Simulation* 19, 4 (2009), 1–20.
- E. F. Fama. 1965. The behavior of stock-market prices. *The Journal of Business* 38, 1 (1965), 34–105.
- E. F. Fama and R. Roll. 1968. Some properties of symmetric stable distributions. *Journal of the American Statistical Association* 63, 323 (1968), 817–836.
- E. F. Fama and R. Roll. 1971. Parameter estimates for symmetric stable distributions. *Journal of the American Statistical Association* 66, 334 (1971), 331–338.
- M. P. Gander and D. A. Stephens. 2007a. Simulation and inference for stochastic volatility models driven by Lévy processes. *Biometrika* 94, 3 (2007), 627–646.
- M. P. Gander and D. A. Stephens. 2007b. Stochastic volatility modelling in continuous time with general marginal distributions: Inference, prediction and model selection. *Journal of Statistical Planning and Inference*.
- J. D. P. Gaver. 1966. Observing stochastic processes, and approximate transform inversion. *Operations Research* 14, 3 (1966), 444–459.
- M. Grabchak et al. 2012. On a new class of tempered stable distributions: moments and regular variation. *Journal of Applied Probability* 49, 4 (2012), 1015–1035.
- M. Hofert. 2011a. Efficiently sampling nested Archimedean copulas. *Computational Statistics & Data Analysis* 55, 1 (2011), 57–70.
- M. Hofert. 2011b. Sampling exponentially tilted stable distributions. *ACM Transactions on Modeling and Computer Simulation* 22, 1 (2011), 1–11.
- D. R. Holt and E. L. Crow. 1973. Tables and graphs of the stable probability density functions. *Journal of Research of the National Bureau of Standards, Section B: Mathematical Sciences* 77 (1973), 143–198.
- P. Hougaard. 1986. Survival models for heterogeneous populations derived from stable distributions. *Biometrika* 73, 2 (1986), 387–396.
- B. Jørgensen. 1987. Exponential dispersion models. *Journal of the Royal Statistical Society. Series B (Methodological)* 49, 2 (1987), 127–162.
- M. Kanter. 1975. Stable densities under change of scale and total variation inequalities. *The Annals of Probability* 3, 4 (1975), 697–707.
- R. Kawai and H. Masuda. 2011. On simulation of tempered stable random variates. *Journal of Computational and Applied Mathematics* 235, 8 (2011), 2873–2887.
- K.-K. Kim and S. Kim. 2016. Simulation of tempered stable Lévy bridges and its applications. *Operations Research* 64, 2 (2016), 495–509.
- U. Küchler and S. Tappe. 2013. Tempered stable distributions and processes. *Stochastic Processes and their Applications* 123, 12 (2013), 4256–4293.
- A. Kyprianou. 2006. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer.
- P. Lévy. 1925. *Calcul des Probabilités*. Gauthier-Villars, Paris.
- B. Mandelbrot. 1960. The Pareto-Lévy law and the distribution of income. *International Economic Review* 1, 2 (1960), 79–106.
- B. Mandelbrot. 1961. Stable Paretian random functions and the multiplicative variation of income. *Econometrica* 29, 4 (1961), 517–543.
- B. Mandelbrot. 1963a. New methods in statistical economics. *Journal of Political Economy* 71, 5 (1963), 421–440.
- B. B. Mandelbrot. 1963b. The variation of certain speculative prices. *The Journal of Business* 36, 4 (1963), 394–419.
- J. H. McCulloch. 1996. Financial applications of stable distributions. In *Statistical Methods in Finance*, Vol. 14 of Handbook of Statistics, pages 393–425. Elsevier
- J. R. Michael, W. R. Schucany, and R. W. Haas. 1976. Generating random variates using transformations with multiple roots. *The American Statistician* 30, 2 (1976), 88–90.

- S. S. Mitra. 1981. Distribution of symmetric stable laws of index 2^{-n} . *The Annals of Probability* 9, 4 (1981), 710–711.
- S. S. Mitra. 1982. Stable laws of index 2^{-n} . *The Annals of Probability* 10, 3 (1982), 857–859.
- E. W. Montroll and J. T. Bendler. 1984. On Lévy (or stable) distributions and the Williams-Watts model of dielectric relaxation. *Journal of Statistical Physics* 34, 1 (1984), 129–162.
- K. A. Penson and K. Górska. 2010. Exact and explicit probability densities for one-sided Lévy stable distributions. *Physical Review Letters* 105 (2010), 210604.
- M. S. Ridout. 2009. Generating random numbers from a distribution specified by its Laplace transform. *Statistics and Computing* 19, 4 (2009), 439–450.
- J. Rosiński. 2001. Series representations of Lévy processes from the perspective of point processes. In *Lévy Processes*, O. Barndorff-Nielsen, S. Resnick, and T. Mikosch (Eds.). Birkhäuser, Boston, pages 401–415.
- J. Rosiński. 2007. Tempering stable processes. *Stochastic Processes and their Applications* 117, 6 (2007), 677–707.
- G. Samoradnitsky and M. S. Taqqu. 1994. *Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance*. CRC Press.
- K.-I. Sato. 1999. *Lévy Processes and Infinitely Divisible Distributions*. Cambridge University Press.
- W. Schoutens. 2003. *Lévy Processes in Finance: Pricing Financial Derivatives*. Wiley.
- H. Stehfest. 1970. Algorithm 368: Numerical inversion of Laplace transforms. *Communications of the ACM* 13, 1 (1970), 47–49.
- V. Todorov et al. 2015. Jump activity estimation for pure-jump semimartingales via self-normalized statistics. *The Annals of Statistics* 43, 4 (2015), 1831–1864.
- M. Tweedie. 1984. An index which distinguishes between some important exponential families. In *Statistics: Applications and New Directions: Proceedings of the Indian Statistical Institute Golden Jubilee International Conference*, pages 579–604.
- V. V. Uchaikin and V. M. Zolotarev. 1999. *Chance and Stability: Stable Distributions and Their Applications*. VSP.
- R. Weron. 1996. On the Chambers-Mallows-Stuck method for simulating skewed stable random variables. *Statistics & Probability Letters* 28, 2 (1996), 165–171.
- V. M. Zolotarev. 1986. *One-dimensional Stable Distributions*. American Mathematical Society.

Received January 2017; revised January 2018; accepted January 2018